

Taxidea taxus jacksoni




Ecosystem-based Automated Range (EBAR)

- Present
- Presence Expected
- Historical

Date Generated: December 5, 2025; Version: 1.0; Stage: Expert Reviewed (Partially Reviewed - National); Scope: Canadian

Synonyms Used: None



0 150 300 km Input Records - 12 GBIF (1934-1993), 5 iDigBio (1948-1991), 5 ON Element Occurrences (1934-2019), 374 ON Observations and Source Features (1895-2019), 3 VertNet (1934-1961); Expert Reviews - Danielle Ethier

Map centre: 80.5154°O 45.4873°N
© NatureServe Canada 2026 under CC BY 4.0

EBAR is relatively coarse scale data and not intended for all applications and analysis. Please see full disclaimer in metadata.

Ecosystem-based Automated Range (EBAR) Metadata

Species

National Scientific Name:	<i>Taxidea taxus jacksoni</i> Schantz, 1946
Scientific Name Reference:	Committee on the Status of Endangered Wildlife in Canada (COSEWIC). 2025. COSEWIC Assessment Results, December 2025. Committee on the Status of Endangered Wildlife in Canada.
National English Name:	American Badger (Southwestern Ontario population)
National French Name:	Blaireau d'Amérique (Population du sud-ouest de l'Ontario); Blaireau d'Amérique de la sous-espèce jacksoni
Element National ID:	741365
Element Global ID:	741362 (go to NatureServe Explorer)
Element Code:	AMAJF04013
Endemism Type:	N
Canadian COSEWIC Name:	
Canadian COSEWIC ID:	621
English COSEWIC Common Name:	American Badger (Southwestern Ontario population)
French COSEWIC Common Name:	Blaireau d'Amérique (Population du sud-ouest de l'Ontario)

Rank/Status

Global Rank:	G5T4 (reviewed April 05, 2016)
National Rank (Canada):	N1 (reviewed 2022)
Subnational Ranks (Canada):	ON=S1
National Rank (United States):	None
Subnational Ranks (United States):	None
National Rank (Mexico):	None
Subnational Ranks (Mexico):	None
Canadian SARA Status:	Endangered/En voie de disparition (June 05, 2003)
Canadian COSEWIC Status:	Endangered
US ESA Status:	None

Range Map

Date Generated:	December 05, 2025
Version:	1.0
Stage:	Expert Reviewed (Partially Reviewed - National)
Scope:	Canadian
Metadata:	Primary Species - <i>Taxidea taxus jacksoni</i> Schantz, 1946; Secondary Species - <i>Taxidea taxus</i> (Schreber, 1777) Input Records - 12 GBIF (1934-1993), 5 iDigBio (1948-1991), 5 ON Element Occurrences (1934-2019), 374 ON Observations and Source Features (1895-2019), 3 VertNet (1934-1961); Expert Reviews - Danielle Ethier
Comments:	Range spatially constrained using a custom polygon to capture the parent species data within the subspecies range extent. Custom polygon based on the 2012 COSEWIC Assessment and Status Report: https://wildlife-species.canada.ca/species-risk-registry/virtual_sara/files/cosewic/sr_blaireau_am_badger_1113_e.pdf
Disclaimer:	Please review our methods document before using EBAR. EBAR range data are relatively coarse scale and appropriate for screening and education purposes, but are not intended for all types of applications and analysis. The absence of data in any geographic areas does not necessarily mean that a species is not present. An ecoshape with a presence value does not necessarily mean that a species is present throughout the entire geographic area.
Presence Definitions:	(Please see Comments above for any exceptions) Present - the species is found within the ecoshape based on species observation data, Element Occurrences, Source Features, Canadian Federal Critical Habitat, or expert opinion. Presence Expected - expert opinion the species may be present, or the ecoshape overlapped with a range estimate or a habitat suitability model. Historical - all species occurrence data within the ecoshape contains observation data greater than 40 years old or an Element Occurrence (EO) that was ranked as Extirpated or Historical (EO Rank of H, H?, X or X?), or expert opinion that the species is extirpated or historical.
Usage Type Definitions:	(Please see Comments above for any exceptions) Breeding - the species is thought to breed within the ecoshape based on eBird Breeding and Behaviour Codes or expert opinion. Possible Breeding - the species is probably or possibly breeding within the ecoshape based on eBird, BBA or jurisdiction Breeding and Behaviour Codes, or on expert opinion.
Map Projection:	North America Albers Equal Area Conic (WKID 4269)

Credits

Suggested Citation:

NatureServe Canada, 2026. Ecosystem-based Automated Range (EBAR) for *Taxidea taxus jacksoni*, Version 1.0, Expert Reviewed (Partially Reviewed - National) (Canadian Scope). Ottawa, Canada. Retrieved from [insert url] on [insert date]

License:

Ecosystem-based Automated Range (EBAR) Project, Copyright NatureServe Canada 2026 under CC BY 4.0 (creativecommons.org/licenses/by/4.0/)

Project Website:

www.natureserve.org/canada/ebar

Contact:

ebbar-kba@natureserve.ca

Input References:

GBIF - [Global Biodiversity Information Facility](https://www.gbif.org/)

iDigBio - [Integrated Digital Biocollection](https://www.idigbio.org/)

ON Element Occurrences - Natural Heritage Information Centre, Ontario Ministry of Natural Resources and Forestry

ON Observations and Source Features - Natural Heritage Information Centre, Ontario Ministry of Natural Resources and Forestry

VertNet - [National Science Foundation](https://www.virgatuscience.org/)

Reviewers by Taxa:

[Reviewers by Taxa](#)