
#### Naemia seriata






#### Ecosystem-based Automated Range (EBAR)

Date Generated: March 26, 2024; Version: 1.0; Stage: Expert Reviewed (National); Scope: Canadian

Synonyms Used: None



Map centre: 64.4208°W 45.0646°N

© NatureServe Canada 2020 under CC BY 4.0

EBAR is relatively coarse scale data and not intended for all applications and analysis. Please see full disclaimer in metadata.



# Ecosystem-based Automated Range (EBAR) Metadata

### Species

| National Scientific Name:  | Naemia seriata (Melsheimer, 1847)   |
|----------------------------|---|
| Scientific Name Reference: | Bousquet, Y., P. Bouchard, A.E. Davies, and D.S. Sikes. 2013. Checklist of beetles (Coleoptera) of Canada and Alaska, second edition. Pensoft Series Faunistica No 109. [Downloadable checklist available: Bousquet, Y., P. Bouchard, A.E. Davies, and D.S. Sikes. 2013. Data associated with Checklist of beetles (Coleoptera) of Canada and Alaska. Second Edition. Data Paper. ZooKeys 360:1-44. http://dx.doi.org/10.5886/998dbs2a] |
| National English Name:     | Seaside Lady Beetle   |
| National French Name:      | Coccinelle côtière  |
| Element National ID:       | 914674  |
| Element Global ID:         | 808504 ( <u>go to NatureServe Explorer</u> )  |
| Element Code:              | IICOLPX030  |
| Endemism Type:             | Ν   |
| Canadian COSEWIC Name:     |   |
| Canadian COSEWIC ID:       |   |
|                            |   |

## Rank/Status

| Global Rank:                       | GNR                |
|------------------------------------|--------------------|
| National Rank (Canada):            | N3 (reviewed 2022) |
| Subnational Ranks (Canada):        | NB=S3, NS=S3       |
| National Rank (United States):     | NNR                |
| Subnational Ranks (United States): | RI=SNR, SC=SNR     |
| National Rank (Mexico):            | None               |
| Subnational Ranks (Mexico):        | None               |
| Canadian SARA Status:              | None               |
| Canadian COSEWIC Status:           | None               |
| US ESA Status:                     | None               |

### Range Map

| Date Generated:         | March 26, 2024   |
|-------------------------|--|
| Version:                | 1.0  |
| Stage:                  | Expert Reviewed (National)   |
| Scope:                  | Canadian   |
| Metadata:               | Primary Species - <i>Naemia seriata</i> (Melsheimer, 1847)<br>Input Records - 92 ACCDC Observations, 51 GBIF, 110 iNaturalist.ca, 76 iNaturalist.org; Expert Reviews - 2 Anonymous   |
| Comments:               | None<br><u>Please see spatial data for Ecoshape-level reviewer comments</u> .  |
| Disclaimer:             | Please review our methods document before using EBAR.  |
|                         | EBAR range data are relatively coarse scale and appropriate for screening and education purposes, but are not intended for all types of applications and analysis.   |
|                         | The absence of data in any geographic areas does not necessarily mean that a species is not present.   |
|                         | An ecoshape with a presence value does not necessarily mean that a species is present throughout the entire geographic area.   |
| Presence Definitions:   | (Please see Comments above for any exceptions)   |
|                         | Present - the species is found within the ecoshape based on species observation data, Element Occurrences, Source<br>Features, Canadian Federal Critical Habitat, or expert opinion.                                       |
|                         | Presence Expected - expert opinion the species may be present, or the ecoshape overlapped with a range estimate or a habitat suitability model.  |
|                         | Historical - all species occurrence data within the ecoshape contains observation data greater than 40 years old or an<br>Element Occurrence (EO) that was ranked as Extirpated or Historical (EO Rank of H, H?, X or X?). |
| Usage Type Definitions: | (Please see Comments above for any exceptions)   |
|                         | Breeding - the species is thought to breed within the ecoshape based on eBird Breeding and Behaviour Codes or expert opinion.  |
|                         | Possible Breeding - the species is probably or possibly breeding within the ecoshape based on eBird, BBA or jurisdiction Breeding and Behaviour Codes, or on expert opinion.   |
| Map Projection:         | North America Albers Equal Area Conic (WKID 4269)  |
|                         |  |

| Credits |
|---------|
|---------|

| Suggested Citation: | NatureServe Canada, 2020. Ecosystem-based Automated Range (EBAR) for Naemia seriata, Version 1.0, Expert Reviewed<br>(National) (Canadian Scope). Ottawa, Canada. Retrieved from [insert url] on [insert date]   |
|---------------------|--|
| License:            | Ecosystem-based Automated Range (EBAR) Project, Copyright NatureServe Canada 2022 under CC BY 4.0<br>( <u>creativecommons.org/licenses/by/4.0/</u> )   |
| Project Website:    | www.natureserve.org/canada/ebar  |
| Contact:            | ebar-kba@natureserve.ca  |
| Input References:   | ACCDC Observations - Atlantic Canada Conservation Data Centre<br>GBIF - <u>Global Biodiversity Information Facility</u><br>iNaturalist.ca - <u>California Academy of Sciences and the National Geographic Society</u><br>iNaturalist.org - <u>California Academy of Sciences and the National Geographic Society</u> |
| Reviewers by Taxa:  | Reviewers by Taxa  |